

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Insecta Mundi

Center for Systematic Entomology, Gainesville,  
Florida

---

10-22-2021

**A new species and significant range extension for the genus  
*Fractipes* Bechyně (Coleoptera: Chrysomelidae: Eumolpinae:  
Eumolpini)**

Wills Flowers

Florida A&M University, rflowers7@earthlink.net

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

---

Flowers, Wills, "A new species and significant range extension for the genus *Fractipes* Bechyně (Coleoptera: Chrysomelidae: Eumolpinae: Eumolpini)" (2021). *Insecta Mundi*. 1381.  
<https://digitalcommons.unl.edu/insectamundi/1381>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

A journal of world insect systematics

# INSECTA MUNDI

---

0890

A new species and significant range extension  
for the genus *Fractipes* Bechyně  
(Coleoptera: Chrysomelidae: Eumolpinae: Eumolpini)

R Wills Flowers  
Center for Biological Control  
Florida A&M University  
Tallahassee, FL 32307, USA

Date of issue: October 22, 2021

Center for Systematic Entomology, Inc., Gainesville, FL

**Flowers RW. 2021.** A new species and significant range extension for the genus *Fractipes* Bechyně (Coleoptera: Chrysomelidae: Eumolpinae: Eumolpini). *Insecta Mundi* 0890: 1–6.

Published on October 22, 2021 by  
**Center for Systematic Entomology, Inc.**  
P.O. Box 141874  
Gainesville, FL 32614-1874 USA  
<http://centerforsystematicentomology.org/>

**INSECTA MUNDI** is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

*Insecta Mundi* is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

**Chief Editor:** David Plotkin, [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Assistant Editor:** Paul E. Skelley, [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Layout Editor:** Robert G. Forsyth

**Editorial Board:** Davide Dal Pos, Oliver Keller, M. J. Paulsen

**Founding Editors:** Ross H. Arnett, Jr., J. H. Frank, Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff

**Review Editors:** Listed on the *Insecta Mundi* webpage

**Printed copies (ISSN 0749-6737) annually deposited in libraries**

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

The Natural History Museum, London, UK

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

**Electronic copies (Online ISSN 1942-1354) in PDF format**

Archived digitally by Portico

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

**Copyright held by the author(s).** This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

# A new species and significant range extension for the genus *Fractipes* Bechyně (Coleoptera: Chrysomelidae: Eumolpinae: Eumolpini)

R Wills Flowers

Center for Biological Control  
Florida A&M University  
Tallahassee, FL 32307, USA  
rflowers7@earthlink.net

**Abstract.** *Fractipes tayrona* **new species** (Coleoptera: Chrysomelidae: Eumolpinae: Eumolpini) is described from a dry forest on the Caribbean coast of Colombia. This locality is very distant from the localities in southeast Brazil, where the other two known species of *Fractipes* Bechyně were described. Male and female genitalia are described for the first time in this genus, and *F. tayrona* is compared with holotype photographs of *F. rhabdopteroides* Bechyně and *F. secundus* Bechyně and Springlová de Bechyně.

**Key words.** Colombia, Brazil, taxonomy, leaf beetle.

**Resumen.** Se describe *Fractipes tayrona* **especie nueva** (Coleoptera: Chrysomelidae: Eumolpinae: Eumolpini) de la costa Caribe de Colombia. La especie fue recolectada en un bosque seco, en un lugar muy lejos del hábitat de los otros dos especies conocidas de *Fractipes* Bechyně, los cuales están conocidas del sureste de Brasil. Se describe los genitales de los machos y hembras por la primera vez en este género, y se compara *F. tayrona* a las fotos de los holotipos de *F. rhabdopteroides* Bechyně and *F. secundus* Bechyně and Springlová de Bechyně.

**Palabras clave.** Colombia, Brasil, taxonomía, escarabajos fitófagos.

**ZooBank registration.** urn:lsid:zoobank.org:pub:6CE2B3FE-4415-4ACA-99D5-8C25FF2CAE3B

## Introduction

Bechyně (1950) described *Fractipes rhabdopteroides* as a new genus and species of eumolpine (Coleoptera: Chrysomelidae: Eumolpinae: Eumolpini) leaf beetle from southeastern Brazil. He described the species as resembling the genus *Rhabdopterus* Lefèvre, except for a very large ventral lobe on the hind femur of the male. Later, Bechyně and Springlová de Bechyně (1961) described a second species, *Fractipes secundus* Bechyně and Bechyně, also from southeastern Brazil, which is distinguished by a very large ventral tooth on the male metafemur. Inspection of a collection of Eumolpinae from a protected area on the Caribbean coast in Colombia has revealed a third species with a very large armament on the male metafemur. This paper describes *Fractipes tayrona* **new species**, and compares it with photos of holotypes of the other two known species. The locality of the new species represents a range extension for the genus of approximately 5,100 km; about as far from the other two species as is possible to get and still be on continental South America.

## Materials and Methods

Specimens examined are in the following institutions:

**FSCA** Florida State Collection of Arthropods, Gainesville, FL, USA

**IAVH** Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, Colombia

**USNM** National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Specimens were examined under a Nikon stereomicroscope. Genitalia dissections were placed in glycerin in polyethylene micro vials and pinned below their respective specimens. Whole specimens were photographed with Auto-Montage software at the Florida State Collection of Arthropods. Genitalia photographs were taken with a Nikon Coolpix 995 through a stereomicroscope, and photos were converted to line drawings with Gimp®. Terminology of the genitalia follows Flowers (1995, 1999), Askevold and Flowers (1994), and Riley and Barney (2015). Label data for holotype, allotype, and paratypes are reproduced verbatim.

## Systematics

### *Fractipes tayrona* Flowers, new species


Figures 1–7, 12–17

**Description.** HOLOTYPE MALE (Fig. 1). Body elongate oval; length 4.7 mm (male range: 4.4–5.3 mm; mean = 4.9 mm;  $n = 6$ ). Upper side strongly shining. Head and pronotum piceous with bronze reflex, elytra dark brown with weak green reflex, labrum reddish brown, maxillary palpi yellow, antennomeres 1–6 and 8–9 reddish brown, 7 and 10–11 darker. Underside dark piceous brown with metallic bronze reflex, apex of sternum VI and sternum




**Figures 1–4.** *Fractipes tayrona*. 1) Holotype male, dorsal habitus. 2) Allotype female, dorsal habitus. 3) Head of male. 4) Underside of male showing femoral spur, and terminal abdominal segments plus pygidium.





**Figures 5–11.** Structures of *Fractipes*. 5) Male hind leg of *F. tayrona*. 6) Male sterna V–VIII of *F. tayrona* with tip of pygidium above. 7) Female sternum VII of *F. tayrona*. 8) Male hind leg of *F. secundus*. 9) Male hind leg of *F. rhabdopteroides* (Figures 8 and 9 are from Bechyně and Springlová de Bechyně [1961]; captions are reversed in the original figure). 10) Holotype and labels of *F. secundus*. 11) Holotype and labels of *F. rhabdopteroides*.



**Figures 12–17.** *Fractipes tayrona*, genitalia. 12) Lateral view of median lobe and partially everted endophallus. 13) En-face view of median lobe. 14) Apical sclerite. 15) En-face view of endophallus at endophallic lateral digits. 16) Spermatheca. 17) Ovipositor.

VII reddish brown. Legs reddish brown, joint of femur–tibia darker. *Head*. Clypeus and frons punctate, punctures separated by distance slightly greater than their diameters. Vertex finely, sparsely punctate with punctures separated by distance greater than their diameters; surface between punctures smooth, shining; antennal calli smooth. Antennomeres 4–6 dorsoventrally compressed and densely setose ventrally (Fig. 3). *Thorax*. Prothorax distinctly wider than long,  $L/W = 0.67$ ; anterior and posterior widths subequal; lateral margins rounded with an obtuse tooth at midlength; anterior and posterior angles prominent, directed laterally. Disc strongly punctate, with punctures separated by a distance equal or slightly less than their own diameters; surface between punctures shining, smooth. Prosternum with scattered pale setae, and with posterior margin of intercoxal process truncate, width of intercoxal process  $0.5\times$  diameter of procoxa. Mesosternum surface alutaceous with scattered pale setae, flat between mesocoxae. Metasternum smooth and shining; metepisternum surface finely alutaceous. *Legs*. Basitarsi of fore and middle legs enlarged, elongate, rectangular. Hind femur with a ventral carina in basal two-thirds, ending in a very large ventral tooth slightly curved inwardly (Fig. 4); carina and apex of tooth black. Apical fourth of hind tibia setose; apex asymmetrically rounded with fine carina on margin (Fig. 5). *Elytra*. Evenly punctate basally, punctate-striate in apical third; punctures separated by distance greater than their diameters; surface between punctures smooth, shining; humeri prominent, subquadrate, width across humeri  $1.3\times$  width across pronotum; basal calli weakly developed; postbasal depression shallow. *Abdomen*. Sterna with numerous short setae and a transverse band of long setae in middle third at apical margins; surface of segments alutaceous. Male sternum VI (Fig. 6) with lateral margins crenulate on apical half, a pair of blunt swellings at posterolateral corners, and a weak depression in center; sternum VII with a deep U-shaped emargination for reception of the apex of the pygidium. Pygidium with a deep longitudinal groove and shallow oval depressions laterally. Terga heavily sclerotized, spicules on terga II and III small and inconspicuous. *Genitalia*. Median lobe in lateral view





**Figure 18.** Dry forest habitat at Parque Nacional Natural Tayrona, Colombia.

(Fig. 12) bent downward at slightly greater than right angle to the basal hood; in en-face view apical margin drawn out into a long acute projection (Fig. 13). Endophallus with a pair of short sclerotized basal lateral digits (Fig. 15) and large, paddle-shaped endophallic lateral digits, sclerotized at their base; a large membranous T-shaped dorsal lobe between the digits, lobe with a pair of lyre-shaped sclerites (Fig. 15). Apical sclerite small, distinctly longer than wide (Fig. 14). Sperm gland elongate and very slender, a very small non-sclerotized cap at the junction of the duct.

**ALLOTYPE FEMALE** (Fig. 2). Body oval; length 5.0 mm (female range: 4.5–5.2 mm; mean = 4.8 mm;  $n = 8$ ); color similar to male but with elytra darker brown. *Head*. With labrum, frons, eyes similar to male; mouthparts similar to male. Antenna filiform, segments 4–6 not flattened and lacking ventral setae. *Thorax*. Prothorax distinctly wider than long,  $L/W = 0.6$ ; shape of pronotum as in male; prosternum similar to male but with setae longer. Meso- and metathoracic sterna as in male. Legs with hind femur slender, unmodified; basitarsi unmodified. *Elytra*. Similar to male but with posthumeral depression deeper. *Abdomen*. Sterna evenly covered with long yellowish setae. Sternum VII with large apical emargination, strongly crenulate laterally (Fig. 7). *Genitalia*. Segments VIII–XI forming very long, thin ovipositor ( $L/W = 24.8$ ; Fig. 17). Sternum VIII with long linear basal apodeme; segment IX covered with minute setae; hemisternites with long basal rods; baculum distinct, apical, slightly shorter than gonocoxae. Gonocoxae longer than wide. Spermatheca (Fig. 16) with receptacle small, scarcely differentiated from the pump.

**Specimens examined.** (6♂♂, 8♀♀). Male HOLOTYPE labeled: COLOMBIA, Magdalena, PNN Tayrona Palangana, 11°20'N 72°2'W, 30 m, Malaise, 05/03–21/03/2001, R. Henriquez Leg. M1483. Female ALLOTYPE labeled: COLOMBIA, Magdalena, PNN Tayrona Palangana, 11°20'N 72°2'W, 30 m, Malaise, 4–17.i.2002, M2756. Holotype and allotype deposited in IAVH. PARATYPES: COLOMBIA, Magdalena: (3♂♂; 2 IAVH, 1 FSCA) same locality, date, and collector as holotype; (1♂, USNM) same locality, date, and collector as allotype; (1♂, 1♀) same locality and collector as holotype, 16.xii.2001–4.i.2002, M2756; (6♀♀; 4 IAVH, 1 FSCA, 1 USNM) PNN Tayrona Gairaca, 11°20'N 74°2'W, 5 m, Malaise, 05/03–21/03/2001, R. Henriquez Leg. M1479.

**Etymology.** This species is named for Parque Nacional Natural Tayrona, where all individuals of this species were collected; name to be treated as a noun.



**Diagnosis.** *Fractipes* differs from all other genera of Neotropical Eumolpinae by the very large size of the male metafemoral projection. *Fractipes tayrona* differs from the two other described species in this genus by the lack of a notch or emargination at the apex of the male hind tibia, by the shape of the projection on the ventral side of the metafemur (Fig. 5, 8–9), and by the dense short setae on the ventral side of male antennomeres 4–6.

## Discussion

In addition to the characters listed above in the diagnosis, *F. tayrona* differs from both *F. rhabdopteroides* (Fig. 9, 11) and *F. secundus* (Fig. 8, 10) in having blunt teeth on the pronotal margins, and a more slender body shape (and consequently a narrower prosternum). All three species share an extreme development in the metafemora of the males, and elongate rectangular basitarsi in the male fore and middle legs. Additionally, *F. tayrona* and *F. rhabdopteroides* share a deeply incised male sternum VII, noted in Bechyně's (1950) description of *F. rhabdopteroides*.

Parque Nacional Natural Tayrona is located on the Caribbean coast of northwestern Colombia. Its dominant vegetation is tropical dry forest (Fig. 18). The photo was taken by the author during the dry season in 2016, 15 years after the specimens were collected and, according to park personnel, after two years of unusually low rainfall. However, the abundant presence of cacti and characteristic dry forest vegetation in the Tayrona forest indicates that *F. tayrona* is well adapted to a seasonally dry climate. This is in contrast to the known habitats of the two Brazilian species, both of which were collected in Brazil's Atlantic Forest.

*Fractipes tayrona* substantially extends the known range of its genus. While the disjunction is striking, it is not unprecedented and perhaps to be expected as more large biodiversity collections are analyzed. For example, a survey of aquatic insects in the Madre de Dios watershed in Peru revealed several mayfly genera formerly considered endemic to southeast Brazil (Sweeney et al. 2009). Even in the older eumolpine literature, there are similar disjunctions, such as Lefèvre's genus *Eurysarcus* in which he included four species from southeast Brazil and one from Manizales, Colombia (Lefèvre 1876, 1878).

## Acknowledgments

I thank Keita Matsumoto and the Natural History Museum, London for the photograph of the holotype of *Fractipes rhabdopteroides*, and Guilherme Ide, Sônia Casari, and Gabriel Biffi of the Museu de Zoologia da Universidade de São Paulo for permission to use the holotype photograph of *Fractipes secundus*. I also thank Michael Geiser and Edward G. Riley for reviewing this manuscript and for their helpful suggestions. Krista E. M. Galley, ELS, of Galley Proofs Editorial Services provided editorial assistance.

## Literature Cited

- Askevold IS, Flowers RW. 1994. *Glyptosceloides dentatus*, a genus and species of Eumolpinae new to Chile (Coleoptera: Chrysomelidae). *Revista Chilena de Entomología* 21: 69–76.
- Bechyně J. 1950. Notes sur les eumolpides de l'Amérique du Sud, (Col. Phytophaga). *Annals and Magazine of Natural History* (Series 12) 3: 70–85.
- Bechyně J, Springlová de Bechyně B. 1961. Insecta Amapaensia: Chrysomeloidea (Col.). *Studia Entomologica* 4: 409–428.
- Flowers RW. 1995. *Hermesia* Lefèvre, a resurrected genus of Neotropical Eumolpinae (Coleoptera: Chrysomelidae). *Proceedings of the Entomological Society of Washington* 97: 35–45.
- Flowers RW. 1999. Internal structure and phylogenetic importance of male genitalia in the Eumolpinae. p. 71–93. In: Cox ML (ed.). *Advances in Chrysomelidae Biology 1*. Backhuys Publishers; Leiden, Netherlands. 691 p.
- Lefèvre E. 1876. Descriptions d'eumolpides nouveaux ou peu connus. *Revue et Magasin de Zoologie (Série 3)* 4: 278–311.
- Lefèvre E. 1878. Voyage de M.E. Steinheil à la Nouvelle Grenade: eumolpides. *Mitteilungen der Münchener Entomologischen Vereins* 2: 112–133.
- Riley EG, Barney RG. 2015. Definition and revision of the *viduatus* species-group of North American *Pachybrachis* Chevrolat (Coleoptera: Chrysomelidae: Cryptocephalinae). *Coleopterists Bulletin* 69: 25–59.
- Sweeney BW, Flowers RW, Funk DH, Ávila S, Jackson JK. 2009. Mayfly communities in two Neotropical lowland forests. *Aquatic Insects* 31(Suppl. 1): 311–318.

Received September 21, 2021; accepted September 29, 2021.

Review editor Michael L. Ferro.